

Cultivation of Medicinal Plants and Development of Plant-Based Pharmaceutical Products under the Arid and Saline Conditions of Karakalpakstan, Uzbekistan

M.K.Kuralbaeva

Independent researcher

e-mail: kuralbayevamamura@mail.com

Abstract: The cultivation of medicinal plants and the development of plant-based pharmaceutical products offer important opportunities for sustainable agriculture and pharmaceutical development in arid regions. The Republic of Karakalpakstan, located in northwestern Uzbekistan, faces major environmental challenges, including soil salinization, water scarcity, drought and land degradation. These conditions create a need for medicinal plant species that can tolerate environmental stress while maintaining adequate biomass production and phytochemical quality. This article reviews the potential of medicinal plant cultivation in Karakalpakstan, with particular emphasis on licorice (*Glycyrrhiza glabra* L.), a medicinal species of pharmaceutical and ecological importance. Published research indicates that licorice can be cultivated on saline soils and that appropriate biological seed treatments may improve seed germination, root biomass and glycyrrhizin content. The article also discusses the relationship between cultivation practices, harvesting, post-harvest processing, phytochemical standardization and the production of plant-based pharmaceutical preparations. A sustainable approach integrating agronomic management, chemical analysis, quality control and conservation of natural plant populations is proposed. Further experimental research is needed to identify cultivation protocols that optimize both pharmaceutical quality and economic viability under the specific environmental conditions of Karakalpakstan.

Keywords: medicinal plants, *Glycyrrhiza glabra* L., Karakalpakstan, soil salinity, medicinal plant cultivation, phytochemical analysis, glycyrrhizin, herbal medicines, pharmaceutical quality.

Introduction

Medicinal plants have long been used as sources of biologically active compounds for traditional medicine and modern pharmaceutical development. Their roots, leaves, flowers, fruits and seeds contain diverse secondary metabolites, including alkaloids, flavonoids, terpenoids, phenolic compounds and saponins. These substances provide valuable starting materials for the development of standardized herbal medicines, pharmaceutical extracts and other plant-derived products.

Despite their potential, medicinal plant production requires careful management because environmental conditions influence plant growth, biomass accumulation and the composition of biologically active compounds. Soil salinity, drought and irregular water availability can substantially affect agricultural productivity and the quality of medicinal raw materials.

These challenges are particularly relevant to Karakalpakstan, where environmental degradation associated with the Aral Sea crisis has contributed to extensive soil salinization and the deterioration of agricultural land. Developing sustainable cultivation systems for medicinal plants could help diversify agricultural production, increase the value of local biological resources and support the development of regional pharmaceutical industries.

Licorice (*Glycyrrhiza glabra* L.) is a promising candidate because of its adaptation to certain saline and arid environments and its importance as a source of glycyrrhizin and other phytochemicals. Research conducted in the Aral Sea region has investigated its cultivation on degraded land and its potential contribution to soil restoration. However, successful production requires an integrated understanding of plant propagation, soil conditions, water management and pharmaceutical quality.

The purpose of this article is to evaluate the scientific basis for cultivating medicinal plants under the arid and saline conditions of Karakalpakstan and to identify practical research priorities for producing high-quality medicinal raw materials and plant-based pharmaceutical products.

Materials and Methods

This article is designed as a narrative review of published scientific literature concerning medicinal plant cultivation, salinity tolerance, phytochemical composition and pharmaceutical processing, with particular attention to research conducted in Uzbekistan and the wider Aral Sea region.

Relevant publications were identified through searches of scientific journal websites and biomedical and academic literature databases. Priority was given to peer-reviewed studies reporting experimental findings on licorice cultivation, biological seed treatments, soil restoration and the chemical characterization of medicinal plant material.

The reviewed evidence was organized into four thematic areas:

Environmental limitations affecting medicinal plant cultivation in Karakalpakstan.

Propagation methods and agronomic management of medicinal plants.

Phytochemical composition and quality assessment of medicinal raw materials.

Processing technologies and quality requirements for plant-based pharmaceutical products.

As this manuscript is a literature-based review, it does not present new experimental measurements or claim to establish original field results. The recommendations are based on published evidence and should be validated through further field and laboratory investigations under local conditions.

Environmental Challenges and Opportunities for Medicinal Plant Cultivation

The principal environmental constraints affecting agricultural production in Karakalpakstan include soil salinity, limited freshwater resources, high temperatures and recurrent drought. These factors influence seed germination, root development, nutrient availability and the accumulation of plant biomass.

High soil salinity can cause osmotic stress and ion imbalance, reducing plant growth and changing metabolic processes. The magnitude of these effects depends on the plant species, developmental stage, soil properties, irrigation regime and duration of exposure to salinity. Consequently, medicinal plants should not be selected solely on the basis of their

traditional medicinal value; their agronomic performance under local environmental conditions must also be evaluated.

Licorice has received particular attention because of its potential to grow in saline environments. Khaitov et al. (2022) discussed its production potential in the harsh environmental conditions of the Aral Sea region, including the possible contribution of licorice cultivation to the rehabilitation of degraded land.

Nevertheless, cultivation should be based on measured soil salinity, groundwater conditions and irrigation-water quality. A species that tolerates moderate salinity may not maintain satisfactory growth or pharmaceutical quality under more severe conditions. Site-specific assessment is therefore essential before establishing commercial plantations.

Propagation and Agronomic Management of Medicinal Plants

The successful cultivation of medicinal plants begins with the selection of suitable propagation methods and high-quality planting material. Depending on the species, propagation may be achieved through seeds, rhizomes, root cuttings or other vegetative structures.

For licorice, seed propagation offers an opportunity to establish plantations on a larger scale. However, seed dormancy, germination variability and unfavorable soil conditions can limit establishment. Appropriate seed preparation, sowing depth, soil management and irrigation practices may improve seedling emergence and subsequent plant development.

An experimental study by Khaitov et al. (2025) investigated biological seed treatments for licorice cultivated on saline land in Karakalpakstan. The experiment was conducted during the 2022 and 2023 growing seasons on soils with electrical conductivity of approximately 10–12 dS m⁻¹. The authors reported that Geogumat treatment increased seed germination by 24.3%, root biomass by 37% and glycyrrhizin content by 12.7% relative to the control. These findings indicate that biological treatments may improve both crop performance and the quality of medicinal raw material under specific saline conditions.

These results should be interpreted within the context of the original experimental conditions. They do not establish that the same treatment will produce identical improvements in every soil type, growing season or cultivation system. Further trials are needed to evaluate reproducibility, application costs, long-term soil effects and the consistency of phytochemical composition.

Water management is another major consideration. Irrigation should meet plant requirements while avoiding excessive salt accumulation in the root zone. Soil electrical conductivity, irrigation-water salinity, drainage and soil moisture should be monitored throughout the growing season. Where feasible, efficient irrigation systems and appropriate soil-management practices may help conserve water and improve cultivation outcomes.

Phytochemical Composition and Pharmaceutical Potential of Licorice

The pharmaceutical value of medicinal plants depends not only on biomass production but also on the identity, concentration and stability of their biologically active compounds.

Licorice roots contain glycyrrhizin, a triterpenoid saponin, together with flavonoids and other secondary metabolites. The composition of the root material can vary according to genetic characteristics, geographical origin, plant age, environmental conditions and cultivation practices.

Hayashi et al. (2003) investigated the chemical characteristics of *Glycyrrhiza glabra* collected in Uzbekistan. Their analysis found that glycyrrhizin content in underground plant parts ranged from 3.3% to 6.1% of dry weight. The study also identified glabridin in the root samples and reported variation in the principal flavonol glycosides detected in leaves. These findings demonstrate the importance of chemical characterization when assessing the pharmaceutical potential of locally sourced licorice.

Environmental stress may affect the production of secondary metabolites, but the relationship is not necessarily linear. An increase in a particular compound does not automatically indicate an improvement in the overall quality or safety of a medicinal

product. Biomass yield, chemical composition, contaminant levels and consistency between batches must be assessed together.

For research in Karakalpakstan, a combined evaluation of root biomass and glycyrrhizin concentration would therefore provide a more meaningful assessment of production potential than biomass measurements alone. Analytical methods such as high-performance liquid chromatography (HPLC) can be used to quantify selected chemical constituents and compare the quality of samples obtained under different cultivation conditions.

From Medicinal Raw Materials to Pharmaceutical Products

The conversion of medicinal plants into pharmaceutical products requires controlled harvesting, post-harvest handling, extraction, analytical testing and appropriate formulation.

The first stage involves identifying the plant correctly and collecting the medicinal part at a suitable developmental stage. For licorice, roots and rhizomes are commonly used as medicinal raw materials. Harvesting procedures should minimize unnecessary damage to the remaining plant population and should be compatible with sustainable production.

After harvesting, the raw material should be cleaned, dried under controlled conditions and stored in a manner that limits moisture uptake, microbial growth and chemical degradation. Drying temperature and duration must be selected according to the characteristics of the plant material and the quality requirements of the intended product.

The principal processing stages may include:

Raw material preparation: botanical identification, cleaning, drying and size reduction.

Extraction: recovery of selected constituents using a suitable solvent and validated extraction procedure.

Concentration and purification: removal of excess solvent and, where necessary, separation of unwanted constituents.

Quality assessment: identification and quantification of marker compounds, determination of moisture content and evaluation of relevant contaminants.

Formulation: development of suitable preparations, such as standardized extracts or other dosage forms, in accordance with applicable pharmaceutical requirements.

For licorice, glycyrrhizin can serve as one of the analytical markers for evaluating extract composition. However, its concentration alone is insufficient to establish the safety, efficacy or overall quality of a finished pharmaceutical product.

Particular attention should be paid to product safety. Glycyrrhizin-containing preparations may cause adverse effects, including elevated blood pressure and potassium imbalance, especially with excessive or prolonged exposure. Therefore, pharmaceutical development must include appropriate dose assessment, safety evaluation, labeling and regulatory compliance.

The production of medicinal products should follow relevant pharmacopoeial standards and good manufacturing practices. Traditional use can help identify promising plants, but it cannot replace chemical characterization, validated production procedures or appropriate safety assessment.

Sustainability and Prospects for Regional Pharmaceutical Development

The development of medicinal plant production in Karakalpakstan has potential economic and environmental benefits. Cultivating suitable species could diversify agricultural activities, create opportunities for local processing and strengthen the supply of plant-derived raw materials.

However, these benefits depend on the sustainable management of natural resources. Uncontrolled harvesting of wild licorice can reduce natural populations and undermine long-term resource availability. Establishing cultivated plantations and implementing responsible harvesting practices may help reduce pressure on wild populations.

A regional development strategy should connect agricultural production with scientific research and pharmaceutical quality control. Cooperation among agricultural researchers, botanists, pharmacognosists, analytical chemists and pharmaceutical manufacturers would support the development of standardized raw materials and reliable processing methods.

Further research should prioritize the following areas:

Screening medicinal plant species for adaptation to different levels of soil salinity and water availability.

Comparing seed and vegetative propagation methods under local environmental conditions.

Evaluating the effects of irrigation, soil amendments and biological seed treatments on biomass and phytochemical composition.

Establishing validated analytical methods for assessing active constituents and contaminants.

Developing standardized extraction procedures and assessing the stability and safety of resulting products.

Conducting economic and environmental assessments of medicinal plant cultivation systems.

These priorities could provide the scientific foundation for integrating sustainable agriculture with pharmaceutical research and production in Karakalpakstan.

Conclusion

The cultivation of medicinal plants under the arid and saline conditions of Karakalpakstan represents a promising area of research at the intersection of sustainable agriculture, pharmacognosy and pharmaceutical technology. Licorice (*Glycyrrhiza glabra* L.) is a particularly relevant candidate because of its medicinal value, potential adaptation to saline environments and capacity to provide roots containing pharmacologically important constituents.

Published studies indicate that appropriate biological seed treatments may improve licorice establishment, root biomass and glycyrrhizin content under certain saline conditions. Nevertheless, successful production requires a broader approach that combines species selection, soil and water management, sustainable harvesting, phytochemical analysis and pharmaceutical quality assurance.

Future investigations should integrate field experiments with laboratory analysis to determine how cultivation conditions affect both the quantity and quality of medicinal raw materials. The resulting evidence could support the development of locally adapted cultivation systems, sustainable use of biological resources and the production of standardized plant-based pharmaceutical products.

In conclusion, strengthening scientific cooperation between agricultural and pharmaceutical research institutions may help transform the medicinal plant resources of Karakalpakstan into a sustainable source of high-quality raw materials and contribute to the development of regional pharmaceutical science.

References:

1. Khaitov, B., Rajabov, N., Murtazayeva, G., Durdiev, N., Norqulov, U., Abdalova, G., Khojasov, A., Rakhmatullaev, Y., Avliyakov, M., Tangirova, G., Mamatkulova, L., Begmatov, I., & Kim, Y. C. (2025). The effectiveness of bio-treatment on licorice (*Glycyrrhiza glabra*) productivity and soil restoration in saline ecosystems. *Eurasian Journal of Soil Science*, 14(3), 198–207. <https://doi.org/10.18393/ejss.1657415>
2. Khaitov, B., Karimov, A., Khaitbaeva, J., Sindarov, O., Karimov, A., & Li, Y. (2022). Perspectives of licorice production in harsh environments of the Aral Sea regions. *International Journal of Environmental Research and Public Health*, 19(18), 11770. <https://doi.org/10.3390/ijerph191811770>
3. Khaitov, B., Tadjetdinov, N., Sayimbetov, A., Khaitbaeva, J., Sindarov, O., Khakberdiev, O., & Nematov, T. (2024). Improving the growth of *Glycyrrhiza glabra* L. in saline soils using bioagent seed treatments. *Eurasian Journal of Soil Science*, 13(1), 43–51.
4. Hayashi, H., Hattori, S., Inoue, K., Khodzhimatov, O., Ashurmetov, O., Ito, M., & Honda, G. (2003). Field survey of *Glycyrrhiza* plants in Central Asia (3). Chemical characterization of *G. glabra* collected in Uzbekistan. *Chemical and Pharmaceutical Bulletin*, 51(11), 1338–1340. <https://doi.org/10.1248/cpb.51.1338>
5. Eisenman, S. W., Zaurov, D. E., & Struwe, L. (Eds.). (2013). *Medicinal Plants of Central Asia: Uzbekistan and Kyrgyzstan*. Springer. <https://doi.org/10.1007/978-1-4614-3912-7>